

★ **Reading: Luke chapter 11. Tong chapter 6**

- Recap: we have discussed spin 0 and spin 1/2 quantum fields. Now move up to spin 1. (Next quarter, we'll discuss renormalizability, and note there the complications with quantizing fields of spin greater than 1.) Examples with spin 1 include non-fundamental (composite) fields, e.g. spin 1 mesons, and also the fundamental force carriers: the photon, gluons, and  $W^\pm$  and  $Z^0$ . The gluons and  $W^\pm$  and  $Z^0$  are associated with non-Abelian groups, which we won't discuss this quarter (we'll see if we get to it next quarter).

- For the massive vector mesons, write down the general lagrangian:

$$\mathcal{L} = -\frac{1}{2}(\partial_\mu A^\nu \partial_\nu A^\mu + a \partial_\mu A^\mu \partial_\nu A^\nu + b A_\mu A^\mu).$$

The sign is chosen so that the kinetic terms of the spatial components have the right sign. Write the EOM and note plane wave solutions  $A_\mu(x) = \epsilon_\nu e^{-ik \cdot x}$  solves it if  $k^2 \epsilon_\nu + a k_\nu (k \cdot \epsilon) + b \epsilon_\nu = 0$ . The longitudinal solutions have  $\epsilon \propto k$  and have mass  $\mu_L^2 = -b/(1+a)$ . The transverse have mass  $\mu_T^2 = -b$ . Can eliminate the uninteresting longitudinal solution by taking  $a = -1$  and  $b \neq 0$ , then write Proca lagrangian

$$\mathcal{L} = -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{1}{2} \mu^2 A_\mu A^\mu,$$

where  $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$ . Each component  $A_\mu$  satisfies the KG equation with mass  $\mu$ . Can choose  $\epsilon^{(\pm)} = \frac{1}{\sqrt{2}}(0, 1, \mp i, 0)$  and  $\epsilon^{(0)} = (0, 0, 0, 1)$ , where the label is the value of  $J_z$  of the spin 1 vector. Normalize by  $\epsilon^{(r)*} \cdot \epsilon^{(s)} = -\delta^{rs}$  and  $\sum_r \epsilon_\mu^{(r)*} \epsilon_\nu^{(r)} = -g_{\mu\nu} + \frac{k_\mu k_\nu}{\mu^2}$ .

The conjugate momenta to  $A_\mu$  are  $\pi^0 = \partial \mathcal{L} / \partial \dot{A}_0 = 0$ , and  $\pi^i = \partial \mathcal{L} / \partial \dot{A}_i = -F^{0i} = E^i$ . Then  $\mathcal{H} = -\frac{1}{2}(F_{0i} F^{0i} - \frac{1}{2} F_{ij} F^{ij} + \mu^2 A_i A^i - \frac{1}{2} \mu^2 A_0 A^0) \geq 0$ .

- Quantize the massive vector:

$$[A_i(t, \vec{x}), F^{j0}(t, \vec{y})] = i \delta_i^j \delta^{(3)}(\vec{x} - \vec{y}).$$

In terms of the plane wave solutions,

$$A_\mu(x) = \sum_{r=1}^3 \int \frac{d^3 k}{(2\pi)^{3/2} (\sqrt{2\omega_k})} \left[ a_k^r \epsilon_\mu^r e^{-ikx} + a_k^{\dagger r} \epsilon_\mu^{*r} e^{ikx} \right],$$

and then

$$[a_k^r, a_{k'}^{\dagger s}] = \delta^{rs} \delta^3(\vec{k} - \vec{k}').$$

and

$$:\mathcal{H} := \sum_r \int d^3k \omega_k a_k^{\dagger r} a_k^r.$$

The propagator, the contraction of  $A_\mu(x)$  and  $A_\nu(y)$ , is

$$\langle T A_\mu(x) A_\nu(y) \rangle = \int \frac{d^4k}{(2\pi)^4} e^{-ik(x-y)} \left[ \frac{-i(g_{\mu\nu} - k_\mu k_\nu / \mu^2)}{k^2 - \mu^2 + i\epsilon} \right].$$

So the Feynman rule is that massive vectors have the momentum space propagator

$$\left[ \frac{-i(g_{\mu\nu} - k_\mu k_\nu / \mu^2)}{k^2 - \mu^2 + i\epsilon} \right].$$

And  $\langle 0 | A_\mu(x) | V(k, r) \rangle = \epsilon_\mu(k)^r e^{-ikx}$ , so incoming vector mesons have  $\epsilon_\mu^r(k)$  and outgoing have  $\epsilon^{*r}(k)$ .