

10/25/19 Lecture outline

• Last time: amplitudes in toy model of real mesons  $\phi$  of mass  $\mu$  and complex nucleons of mass  $m$ , with  $H_{int} = -g\phi\bar{\psi}\psi$ . Get

$$\mathcal{A}_{NN \rightarrow NN} = (-ig)^2 \left[ \frac{1}{(p_1 - p'_1)^2 - \mu^2 + i\epsilon} + \frac{1}{(p_1 - p'_2)^2 - \mu^2 + i\epsilon} \right] (2\pi)^4 \delta^{(4)}(p_1 + p_2 - p'_1 - p'_2) + \mathcal{O}(g^4).$$

Explicitly, in the CM frame,  $p_1 = (\sqrt{p^2 + m^2}, p\hat{e})$  and  $p_2 = (\sqrt{p^2 + m^2}, -p\hat{e})$ ,  $p'_1 = (\sqrt{p^2 + m^2}, p\hat{e}')$ ,  $p'_2 = (\sqrt{p^2 + m^2}, -p\hat{e}')$ , where  $\hat{e} \cdot \hat{e}' = \cos\theta$ , and get

$$\mathcal{A} = g^2 \left( \frac{1}{2p^2(1 - \cos\theta) + \mu^2} + \frac{1}{2p^2(1 + \cos\theta) + \mu^2} \right).$$

Scattering by  $\phi$  exchange leads to an attractive Yukawa potential. This was Yukawa's original goal, to explain the attraction between nucleons. Indeed, the t-channel term in e.g. the above  $N+N$  scattering amplitude gives, upon using  $(p_1 - p'_1)^2 - \mu^2 = -(|\vec{p}_1 - \vec{p}'_1|^2 + \mu^2)$ , and the Born approximation<sup>1</sup> in NRQM,  $\mathcal{A}_{NR} = \int d^3\vec{r} e^{-i(\vec{p}' - \vec{p}) \cdot \vec{r}} V(\vec{r})$ , the attractive Yukawa potential

$$V(r) = \int \frac{d^3q}{(2\pi)^3} \frac{-(g/2m)^2 e^{i\vec{q} \cdot \vec{r}}}{|\vec{q}|^2 + \mu^2} = -\frac{(g/2m)^2}{4\pi r} e^{-\mu r}.$$

(The  $1/(2m)^2$  is because we normalized the relativistic states with the extra factor of  $2E \approx 2m$  as compared with standard nonrelativistic normalization<sup>2</sup>. This gives Yukawa's explanation of the attraction between nucleons, from meson exchange. The u-channel term is an exchange potential interaction, which exchanges the positions of the two identical particles in addition to giving a potential. For angular momentum  $\ell$  in a partial-wave expansion, the exchange term differs from the direct one by a factor of  $(-1)^\ell$ .

• More examples:

(1)  $N(p_1) + \bar{N}(p_2) \rightarrow N(p'_1) + \bar{N}(p'_2)$  has

$$i\mathcal{A} = (-ig)^2 \left( \frac{i}{(p_1 - p'_1) - \mu^2} + \frac{i}{(p_1 + p_2) - \mu^2} \right).$$

<sup>1</sup> Max Born, in QM, or Lord Rayleigh classically:  $\frac{d\sigma}{d\Omega} \sim |U(\vec{q})|^2$ .  $S - 1 \approx V$  so  $\langle \vec{p}', -\vec{p}' | S - 1 | \vec{p}, -\vec{p} \rangle \approx \langle \vec{p}', -\vec{p}' | V | \vec{p}, -\vec{p} \rangle$

<sup>2</sup> This is clear on dimensional grounds, since  $[g] \sim m$ . Write  $\langle f | U | i \rangle / \sqrt{\langle f | f \rangle \langle i | i \rangle}$  for the properly normalized amplitude. More generally, write  $a(p) = \sqrt{2E} \hat{a}(p)$  and  $\mathcal{A} = \prod_i \sqrt{2E_i} \prod_f \sqrt{2E_f} \hat{\mathcal{A}}$ .

(2)  $N(p_1) + \bar{N}(p_2) \rightarrow \phi(p'_1)\phi(p'_2)$  has

$$i\mathcal{A} = (-ig)^2 \left( \frac{i}{(p_1 - p'_1) - m^2} + \frac{i}{(p_1 - p'_2) - m^2} \right).$$

(3)  $N(p_1) + \phi(p_2) \rightarrow N(p'_1) + \phi(p'_2)$  has

$$i\mathcal{A} = (-ig)^2 \left( \frac{i}{(p_1 - p'_2) - m^2} + \frac{i}{(p_1 + p_2) - m^2} \right).$$

Note: the  $1/2!$  from expanding  $e^{-i \int d^4x \mathcal{H}_I(x)}$  is cancelled by a factor of 2 from exchanging the two vertices.

- Crossing symmetry, CPT. Write  $1 + 2 \rightarrow \bar{3} + \bar{4}$ . Take all momenta incoming,  $\mathcal{A}(p_1, p_2, p_3, p_4)$ , with  $p_1 + p_2 + p_3 + p_4 = 0$  and use  $s = (p_1 + p_2)^2$ ,  $t = (p_1 + p_3)^2$  and  $u = (p_1 + p_4)^2$ . Note  $s + t + u = \sum_{n=1}^4 m_n^2$ . The process  $1 + 2 \rightarrow \bar{3} + \bar{4}$  is kinematically allowed for  $s > 4m^2$ ,  $t < 0$ ,  $u < 0$ . If instead  $u > 4m^2$ , it's the process  $1 + 3 \rightarrow \bar{2} + \bar{4}$ .

- We saw above that the  $t$  channel term above is associated with the Yukawa potential. The  $u$  channel term is similar. Now consider the  $s$  channel, in e.g. the  $N + \bar{N}$  scattering amplitude. Using the CM relations  $\vec{p}_1 = -\vec{p}_2 \equiv \vec{p}$  and  $E_1 = E_2 = \sqrt{p^2 + m^2}$  gives

$$\mathcal{A} \sim \frac{1}{4m^2 + 4p^2 - \mu^2 + i\epsilon},$$

so for  $\mu < 2m$  the denominator is always positive, and the amplitude decreases with increasing  $p^2$ . For  $\mu > 2m$  there is a pole at  $(p_1 + p_2)^2 = \mu^2$ , where the intermediate meson goes on shell. This leads to a peak (not a pole, of course; because the intermediate particle is unstable anyway, the denominator gets an imaginary contribution from higher order contributions), a *resonance*, in the cross section. E.g.  $Z_0$  pole in  $e^+e^- \rightarrow \mu^+\mu^-$ , but not in  $e^+e^- \rightarrow \gamma\gamma$ .

- Solve  $\mathcal{L} = \frac{1}{2}\partial\phi^2 - \frac{1}{2}m^2\phi^2 - J(x)\phi$ . Using Dyson + Wick's theorem,  $U(\infty, -\infty) =: e^{O_1 + \frac{1}{2}O_2}$  ;, where  $O_1 = -i \int d^4x J(x)\phi(x)$  and  $O_2 = (-i)^2 \int d^4x_1 d^4x_2 D_F(x_1 - x_2) J(x_1) J(x_2)$ . So  $O_2 = \alpha + i\beta$  is a number, whereas  $O_1$  is an operator. Will lead to probability  $P_n$  for creating out of the vacuum a state with  $n$  mesons given by  $P_n = e^{-|\alpha|} |\alpha|^n / n!$ , the Poisson distribution. You'll work out the details in the HW assignment.

- Compute probabilities by squaring the S-matrix amplitudes, but have to be careful with the delta functions, since squaring the delta functions would give nonsense.

Warmup: consider quantum mechanics, with  $U(t) = T e^{-i \int^t H(t) dt}$ ,

$$\langle f|U(t)|i \rangle \approx -i \langle f|H_{int}|i \rangle \int_0^t dt e^{i\omega t},$$

where  $\omega = E_f - E_i$ . If we take  $t \rightarrow \infty$  first, we get  $\delta(\omega)$  and squaring would give nonsense. That's because we're asking the wrong question if we ask about probability for a transition over all time – instead, we should ask about the rate. So keep  $t$  finite for now. Squaring gives  $P(t) = 2|\langle f|H_{int}|i\rangle|^2(1 - \cos\omega t)/\omega^2$ . For  $t \rightarrow \infty$ , multiply by  $dE_f\rho(E_f)$  and replace  $(1 - \cos\omega t)/\omega^2 = 4\sin^2(\frac{1}{2}\omega t)/\omega^2 \rightarrow \pi t\delta(\omega)$  (using  $\int_{-\infty}^{\infty} dx x^{-2} \sin^2 x = \pi$  (hint:  $\sin^2 x/x^2 = (2 - e^{i2x} - e^{-i2x})/4x^2$  and close the contour in the correct directions)) to get

$$\dot{P}_{i \rightarrow f} = 2\pi|\langle f|H_{int}|i\rangle|^2\rho(E).$$

This is “Fermi’s Golden Rule” – it was actually derived by Dirac, but Fermi used it a lot and called it the golden rule. Another aside: Fermi and Dirac independently discovered that spin 1/2 objects must anticommute, and Dirac generously named such objects “Fermions”.

Naively taking  $t \rightarrow \infty$  initially would have given amplitude  $\sim \delta(\omega)$ , and squaring that would give  $\delta(\omega)^2$ , which needs to be replaced with  $\delta(\omega)2\pi T$ , and then divide by  $T$  to get the rate. Similarly in field theory,  $\delta(p)^2$  should be replaced with probability  $\sim \delta(p)$  times phase space volume factors.