

- Continue with lecture 3 mathematica notebook examples (on TED side), of examples of residues, poles, verifying that the CR equations are satisfied, and noting that they imply that $\Re f(z)$ and $\Im f(z)$ are solutions of the 2d Laplace equations; examples. Useful for e.g. some electrostatics problems. Verify it for some examples using Mathematica.

- Discuss $\oint_C dz/z = 2\pi i$ in terms of $z^{-1} = \partial_z \log z$ and the behavior of $\log z$ in the complex plane. Write $f(z)dz$ in terms of real and imaginary parts, and then as $(\vec{F} \cdot d\vec{\ell}, (d\vec{\ell} \times \vec{F}))$, with $\vec{F} = (u, -v)$, and note that the CR equations imply that \vec{F} has no divergence or curl, clarifying why $\oint f(z)dz$ is “almost zero”, up to the effects from the poles. Indeed, the poles are places where singularities of the derivatives of a certain type. This is related to the fact that $\log(z - z_0)$ is a Green’s function for the 2d Laplacian. We will discuss Green’s functions later.

- Recall the example of $\int_{-\infty}^{\infty} dx(1 + x^2)^{-1} = \pi$, and show that one gets the same answer if C is closed instead in the lower half plane, accounting for the sign convention.

- Other examples of evaluating integrals by Cauchy’s theorem. $\int_0^\pi d\theta/(a + b \cos \theta) = \pi/\sqrt{a^2 - b^2}$,

- Residues and poles of $\pi/\sin(\pi z)$ and $\pi \cos(\pi z)/\sin(\pi z)$ and applications of Cauchy’s theorem to evaluate some sums, $\sum_{n=1}^{\infty} f(n)$. Examples in the mathematica notebook.

- Example: consider an L, R circuit, driven by source $V(t) = A \int e^{i\omega t} d\omega/2\pi$; this source corresponds to a voltage spike at time $t = 0$. Find $I(t) = A \int (R + i\omega L)^{-1} d\omega/2\pi$. Discuss where to close the contour and get $I(t < 0) = 0$ and $I(t > 0) = (A/L)e^{-Rt/L}$. Makes sense.

Ended here. Continue with below next time

- Gamma function $\Gamma(z)$; give integral definition and type it into mathematica, $\Gamma(z + 1) = z\Gamma(z)$ and relation to factorial. Poles at $x = 0$ and negative integers. Check with mathematica. Also $\Gamma(z)\Gamma(1 - z) = \pi/\sin(\pi z)$.

- Gaussian integral, including in multi-dimensions. Normalization of the normal distribution. Relation to spherical integrals and solid angles.

- Example of forced SHO with $F(t) = F \cos(\omega t)$ and the particular solution for both $\omega \neq \omega_0$ and $\omega = \omega_0$.

- Suppose that we want to solve the ODE $\frac{d^2 x}{dt^2} = f(x, \dot{x})$, where f is some given function, e.g. $f = -\omega_0^2 x - \gamma v$ for the case of a damped SHO. Note that we are here taking $f(x, \dot{x})$ to not depend explicitly on t . Plot (x, \dot{x}, t) curve, so $(dx, dv, dt) = dt(v, f, 1)$ is the tangent vector. Project to the (x, v) plane, and plot (v, f) to give the tangent vector field.

For example, for the SHO, $f = -x$ so the tangent vector field in the (x, v) plane is $(v, -x)$, i.e. the phase space motion is a circle. Discuss example in cell 1.6 of Chapter1.nb. It is often useful to use p instead of \dot{x} (in simple cases, this is just a rescaling as $p = m\dot{x}$). Plot phase space motion for the solution of the undamped SHO vs the damped SHO.

- Non- dissipative systems have conserved energy and the flow in the (x, v) plane has zero divergence. Hence the area in phase space is constant in time.

- Hamiltonian flows: $H(x, p, t)$ with $\dot{x} = \partial_p H$ and $\dot{p} = -\partial_x H$. Discuss \dot{H} vs $\partial_t H$ and show that $\dot{H} = 0$ if $\partial_t H = 0$: this is conservation of energy if the system does not explicitly depend on t . You will learn more about this in physics 110.

- Following Dubin 1.4.3, discuss Euler's method for numerical solutions of differential equations. First consider $\dot{f} = f(t, v)$ e.g. for $f(t, v) = t - v$.