

2/1/16 Homework 3. Due Feb 9

1. As mentioned in lecture, the Coleman-Weinberg potential for  $V_{int} = \frac{\lambda}{4!}\phi^4$  is

$$\begin{aligned} V_1(\bar{\phi}) &= i \sum_{n=1}^{\infty} \frac{1}{2n} \int \frac{d^4k}{(2\pi)^4} \left( \lambda \frac{1}{k^2 - m^2 + i\epsilon} \frac{\bar{\phi}^2}{2} \right)^n \\ &= \frac{1}{2} \int \frac{d^4k_E}{(2\pi)^4} \ln \left( 1 + \frac{\frac{1}{2}\lambda\bar{\phi}^2}{k_E^2 + m^2} \right) \end{aligned}$$

(S. Coleman and E. Weinberg.) Symmetry factors:  $1/n!$  not all the way cancelled, because of  $Z_n$  rotation symmetry, and reflection, gives  $1/2n$ . At each vertex, can exchange external lines, so  $1/4!$  not all the way cancelled, leads to  $1/2$  for each vertex. Here you will derive the last expression above for  $V_1(\bar{\phi})$  another way. Write

$$e^{iW[J]} = \int [d\phi] e^{i(S + \int J\phi)/\hbar}$$

and expand  $\phi(x) = \bar{\phi} + \eta(x)$ , treating  $\bar{\phi}$  as a constant and imagining  $\eta$  to be a small fluctuation, and keeping only terms to order  $\eta^2$ . Do the Gaussian integral over  $\eta$  formally. Then use the relation given in class to convert  $W[J]$  into  $\Gamma[\bar{\phi}]$  to finally reproduce the above expression for  $V_1(\bar{\phi})$ . Hint: use  $\ln \det B = \text{Tr} \ln B$  for any operator  $B$ , so e.g.  $\ln \det(\partial^2 + m^2) = V_4 \int \frac{d^4k}{(2\pi)^4} \ln(-k^2 + m^2)$  (where  $V_4$  is a spacetime box size which can be taken to infinity at the end of the day; it cancels anyway).

- 2 Consider  $\mathcal{L} = \partial_\mu \phi^* \partial^\mu \phi - m^2 \phi^* \phi - \frac{\lambda}{4} \phi^2 \phi^{*2}$ , where  $\phi$  is a complex scalar field. The lagrangian has a symmetry,  $\phi \rightarrow e^{ia} \phi$ , so there is a corresponding conserved current  $J^\mu(x)$ , with  $\partial_\mu J^\mu = 0$ . Find  $J^\mu(x)$ , using the Noether method. Now, in analogy with the above problem, use the functional integral to show that the current is conserved in correlation functions, up to contact terms:

$$\frac{\partial}{\partial x} \langle T J^\mu(x) \phi(y) \rangle = \beta \delta^4(x - y) \langle \phi(0) \rangle.$$

Derive this similarly to problem 2 of HW 2, and determine the constant  $\beta$ .

3. Consider a diagram, which is a two-loop contribution to the propagator in  $\phi^4$  theory, which looks like a propagator that goes through the middle of a circle. Compute the diagram in the limit of zero mass for  $\phi$ , using dim reg. Show that for  $\epsilon = 4 - d \ll 1$  it takes the form

$$-ip^2 \frac{\lambda^2}{12(4\pi)^4} \left[ -\frac{1}{\epsilon} + \log p^2 + \dots \right].$$

The coefficient involves a Feynman parameter integral that can be evaluated for  $d = 4$ .